



2.6F PERIPHERALLY INSERTED CENTRAL CATHETER (PICC)

VASCU-PICC® TLS

Catalog Code	Description	QTY
MR17012608	2.6F X 50CM DUAL LUMEN PICC WITH CUFF	5/BOX

Basic Set Contains:

- | | |
|-----------------------------------|---------------------------|
| (1) Catheter with Cuff and Stylet | (2) Needleless Connectors |
| (1) Adaptor w/Sideport | (1) Tourniquet |
| (1) 3F OTN Tearaway Introducer | (1) Tape Measure |
| (3) Gauze | (1) Securement Device |
| (1) Syringe | (1) Patient Chart Sticker |

IMPORTANT RISK INFORMATION

Indications For Use: 2.6F Peripherally Inserted Central Vein Access Catheters are indicated for short or long term access to the central venous system via peripheral insertion in neonates, infants, and children. It may be used for administration of fluids, medication, and nutritional therapy.

Contraindications:

- This catheter is not intended for any use other than that which is indicated. Do not implant catheter in thrombosed vessels.
- The presence of skin related problems around the insertion site (infection, phlebitis, scars, etc.)
- The presence of device related bacteremia or septicemia.
- Previous history of venous/subclavian thrombosis or vascular surgical procedures at insertion site.
- Fever of unknown origin.
- The patient's body size is insufficient to accommodate the size of the implanted device.
- The patient is known or is suspected to be allergic to materials contained in the device.
- Past irradiation of prospective insertion site.
- Local tissue factors will prevent proper device stabilization and/or access.



2.6F PERIPHERALLY INSERTED CENTRAL CATHETER (PICC)

VASCU-PICC® TLS



VASCU-PICC® TLS

Approved for infusion therapy

VASCU-PICC® TLS is the latest addition to our PICC product line and features a dual lumen configuration and a soft polyurethane body. Since many pediatric patients require a combination of nutrition, fluids and/or medicine, the dual lumen configuration enables one catheter to serve multiple functions.

